

CHAPTER 49

Coronary Artery Disease

KEY TEACHING POINTS

- In the evaluation of patients with chronic intermittent chest pain and suspected coronary artery disease, the most helpful bedside finding is the patient's description of pain (i.e., typical angina, atypical angina, or nonanginal chest pain). In these patients the following findings increase probability of coronary disease, but only modestly: an ankle-to-arm pressure index of 0.9 or less, arcus senilis, and the earlobe crease.
- In the evaluation of patients with sustained chest pain and suspected myocardial infarction, the most helpful bedside finding is the electrocardiogram. In these patients the following findings increase probability of myocardial infarction: systolic blood pressure less than 100 mm Hg, the third heart sound, jugular venous distention, diaphoresis, and crackles.
- The finding of chest wall tenderness *decreases* probability of myocardial infarction in patients with sustained chest discomfort being evaluated in the emergency department, but it is unhelpful when evaluating chronic intermittent chest pain in the clinic.
- In patients with suspected myocardial infarction, the response to nitroglycerin and to GI cocktail is unhelpful.
- In patients with suspected acute coronary syndromes, the following combination of findings identifies a group of patients at very low risk for complications in the next 24 hours: an electrocardiogram without ST/T wave changes, pain severity that is less severe than prior angina, the absence of hypotension, and the absence of crackles.

I. INTRODUCTION

Coronary disease is the leading cause of heart disease and death in the United States,¹ and chest pain accounts for 8% to 10% of complaints of patients presenting to clinics or emergency departments.²⁻⁴ The bedside diagnosis of chest pain is difficult and at times humbling, as illustrated by up to 1% to 8% of patients with myocardial infarction (confirmed by cardiac biomarkers) being misdiagnosed and discharged home from emergency departments.⁵⁻¹² The focus of this chapter is to identify all aspects of the initial patient encounter—patient interview, physical examination, and the electrocardiogram—that help to distinguish patients with angina and myocardial infarction from those with mimicking disorders.

The first clear description of **angina pectoris** was given in 1768 by William Heberden, who coined the term* and provided a clinical description that has been unsurpassed. Just 8 years later, Edward Jenner linked angina to “ossification” of the coronary arteries and insufficient coronary blood flow,¹³ and in 1878 (more than 50 years before the introduction of electrocardiography), Adam Hammer correctly diagnosed the first case of myocardial infarction during life in a young man with sudden collapse, bradycardia, and enfeebled heart tones.^{14,15} Coronary disease was once considered to be an uncommon disorder—the great 19th century American cardiologist Austin Flint found only seven cases of angina in his clinical records¹⁶ and Osler personally observed only 40 cases during his career.¹³

II. THE FINDINGS

A. INTRODUCTION

Unlike other clinical problems in cardiology, such as valvular disease and heart failure, patients with coronary artery disease have few or no physical findings. For more than 100 years, the most important aspect of diagnosing coronary disease has been the patient’s description of chest pain, whereas the most important element in diagnosing myocardial infarction (at least since 1918) has been the electrocardiogram.

B. DESCRIPTION OF CHEST PAIN

Heberden wrote that **angina** is a “most disagreeable sensation in the breast” that seizes patients “while they are walking” yet vanishes “the moment they stand still.”¹⁷ Modern definitions of **typical angina** retain most of Heberden’s essential features, by defining it as substernal discomfort with three characteristics: it (1) is precipitated by exertion, (2) is improved by rest or nitroglycerin (or both), and (3) lasts less than 10 minutes. Many patients also describe radiation of the pain to the shoulders, jaw, or inner aspect of the arm. In contrast, **atypical angina** is substernal discomfort with atypical features (e.g., it is not always relieved by nitroglycerin, not always brought on by exertion, or relieved after 15 to 20 minutes of rest), and **non-anginal chest pain** lacks all features of typical angina (i.e., it is unrelated to activity, unrelieved by nitroglycerin, or otherwise not suggestive of angina).

C. HAND GESTURES DURING DESCRIPTION OF CHEST PAIN

According to traditional teachings, patients provide diagnostic clues to the physician by the hand gestures they spontaneously make when describing their chest pain. Four of these gestures are: (1) **Levine sign**—placing clenched fist against the sternum, (2) **palm sign**—placing the extended palm against the sternum, (3) **arm sign**—gripping the left arm, and (4) **pointing sign**—pointing to a single point on the chest with one or two fingers.¹⁸ According to traditional teachings, gestures suggesting deep, poorly localized visceral pain (Levine and palm signs) or pain radiating to the left arm (arm sign) increase the probability of coronary disease, whereas gestures indicating well-localized somatic pain (pointing sign) decrease the probability of disease.

* Heberden based the term *angina* on the Greek *agkhone*, which means “strangling.” This Greek root also forms the basis for the English words *anxiety* and *anguish*. Heberden’s selection of *angina* was unfortunate because the term had already been applied to other conditions of the throat, such as Vincent *angina* and Ludwig *angina*.

D. PHYSICAL FINDINGS

Some of the findings that appear in [EBM Boxes 49.1 and 49.2](#) are discussed in other chapters: crackles ([Chapter 30](#)), displaced precordial pulsation ([Chapter 38](#)), and the third heart sound ([Chapter 41](#)).

I. EARLOBE CREASE

The earlobe crease is a diagonal crease across the earlobe, connecting the lowest point on the tragus to the outside of the earlobe ([Fig. 49.1](#)). Some investigators define the finding as a crease traversing at least one-third the distance from tragus to posterior pinna,^{37,38} whereas others require the crease to extend the total distance.^{29,32,39} In a letter to the editor written in 1973⁷² Frank first presented the “positive earlobe sign” as a sign tightly associated with other cardiovascular risk factors. Although its association with coronary disease remains controversial and its pathogenesis a mystery, many investigators have shown that the earlobe crease is a modest risk factor for coronary artery disease, independent of other traditional risk factors, such as hypertension, age, diabetes mellitus, family history, hyperlipidemia, obesity, and cigarette smoking.^{32,37,39,73,74}

2. ARCUS SENILIS

Arcus senilis is a white or grayish opaque ring about the circumference of the cornea. Since the 1830s this sign has been associated with both older age (hence “senilis”) and vascular disease (Virchow considered it a definite sign of heart disease).⁷⁵ Modern investigators^{76,77} continue to suggest arcus senilis is linked to coronary disease, independent of its association with hyperlipidemia, although others challenge this view.⁷⁵

3. ANKLE-TO-ARM PRESSURE INDEX

After positioning the patient supine, the clinician uses a handheld Doppler stethoscope to measure the highest systolic blood pressure in the posterior tibial or dorsalis pedis artery (i.e., the “ankle” pressure). The **ankle-to-arm pressure index** represents this ankle pressure divided by the systolic pressure in the brachial artery (see [Chapter 54](#)).

E. GI COCKTAIL

For many years, clinicians working in emergency departments have mixed liquid antacids with other substances (most commonly viscous lidocaine, a topical anesthetic, and an elixir with the trade name of Donnato!, an antispasmodic) to create **GI cocktails**, which are administered orally to patients presenting with chest or upper abdominal discomfort. Because GI cocktail should act topically only on GI mucosa, prompt relief of a patient’s discomfort is said to support a GI cause of pain (and, by inference, argue against a cardiac cause of the pain). Although antacid, lidocaine, and Donnato! are the standard ingredients of GI cocktail, some investigators have shown that antacid alone (without lidocaine or Donnato!) may relieve pain just as well.⁷⁸

III. CLINICAL SIGNIFICANCE

A. DIAGNOSING CORONARY ARTERY DISEASE

[EBM Box 49.1](#) summarizes the accuracy of bedside findings in diagnosing coronary artery disease (based on study of more than 10,000 patients).⁷⁹ Almost all of the patients in these studies presented to outpatient clinics with intermittent chest pain, and the diagnosis of coronary artery disease was based on subsequent cardiac




EBM BOX 49.1
*Coronary Artery Disease**

| Finding (Reference) [†] | Sensitivity (%) | Specificity (%) | Likelihood Ratio [‡] if Finding Is | |
|---|--------------------|--------------------|--|------------|
| | | | Present | Absent |
| Patient Interview | | | | |
| Description of Chest Pain | | | | |
| Classification of chest pain ¹⁹⁻²⁶ | | | | |
| Typical angina | 50-91 | 78-94 | 5.8 | — |
| Atypical angina | 8-44 | — | 1.2 | — |
| Nonanginal chest pain | 4-22 | 14-50 | 0.1 | — |
| Pain duration >30 min ²⁷ | 1 | 86 | 0.1 | NS |
| Associated dysphagia ²⁷ | 5 | 80 | 0.2 | NS |
| Other | | | | |
| Male sex ^{24,25,28-33} | 72-86 | 36-58 | 1.7 | 0.3 |
| Age ^{25,28-30,33,34} | | | | |
| <30 years | 0-1 | 97-98 | NS | — |
| 30-49 year | 16-38 | — | 0.6 | — |
| 50-70 years | 62-73 | — | 1.3 | — |
| >70 years | 2-52 | 67-99 | 2.6 | — |
| Prior myocardial infarction ^{24,26,30,31,33,35,36} | 42-69 | 66-99 | 3.8 | 0.6 |
| Physical Examination | | | | |
| Earlobe crease ^{29,32,37-41} | 26-80 | 33-96 | 2.3 | 0.5 |
| Arcus senilis ⁴² | 40 | 86 | 3.0 | 0.7 |
| Chest wall tenderness ^{27,43-45} | 1-69 | 16-97 | 0.8 | NS |
| Ankle-to-arm pressure index <0.9 ^{46,47} | 20-26 | 93-95 | 4.0 | 0.8 |
| Laterally displaced apical impulse ⁴⁸ | 5 | 100 | NS | NS |
| Electrocardiogram | | | | |
| Normal ^{26,48,49} | 15-33 | 50-69 | NS | NS |
| ST/T wave abnormalities ^{19,26,36} | 14-44 | 73-93 | NS | NS |

*Diagnostic standard: for *coronary artery disease*, coronary angiography reveals greater than 50%,^{20,22-24,26,28,29,32,34,35,37-40,42,44,46,47,49} >60%,⁴⁸ or >70% to 75%^{19,21,25,27,30,31,33,36,41} stenosis of any epicardial vessel or positive myocardial perfusion scan.⁴⁵
[†]Definition of findings: for classification of chest pain, earlobe crease, and arcus senilis, see the text.
[‡]Likelihood ratio (LR) if finding present = positive LR; LR if finding absent = negative LR.
 NS, Not significant.

[Click here to access calculator](#)



catheterization revealing a significant stenosis (>50% to 70% luminal narrowing) in any major epicardial vessel (i.e., single-vessel disease or worse).

According to the likelihood ratios (LRs) in [EBM Box 49.1](#) the findings *increasing* the probability of coronary disease the most in patients with intermittent chest pain are typical angina (LR = 5.8), ankle-to-arm pressure index of less than 0.9 (LR = 4), previous myocardial infarction (LR = 3.8), arcus senilis (LR = 3), age older than 70 years (LR = 2.6), and a positive earlobe crease (LR = 2.3).

These studies confirm Heberden's original impression that the key diagnostic finding in patients with chest pain is the patient's actual description of pain. Many investigators have attempted to improve on Heberden's definition of typical angina by dissecting apart the individual components of the patient's description (e.g., response to nitroglycerin or the pain's quality) or by creating complicated angina scoring schemes, but each of these attempts to improve diagnosis is less accurate than the clinician's global perception of whether the patient's pain is typical angina or not.⁷⁹

The findings that *decrease* the probability of coronary artery disease in these studies are chest pain that is nonanginal (i.e., pain unrelated to activity, unrelieved by nitroglycerin, or otherwise not suggestive of angina, LR = 0.1), pain duration longer than 30 minutes (LR = 0.1), and associated dysphagia (LR = 0.2).

Unhelpful findings include atypical angina, chest wall tenderness, and a displaced apical impulse. Additional descriptors of the pain, such as burning pain, pain made worse by food or emotion, and radiation of the pain to the arms, are also unhelpful (i.e., they appear just as often in patients with coronary disease as in patients with noncardiac chest pain, and the LRs are not different from the value of 1).⁷⁹ Neither the Levine sign nor the palm sign affects the probability of coronary disease.⁸⁰ Interestingly, electrocardiographic findings (i.e., normal vs. abnormal, presence or absence of nonspecific ST changes) also are diagnostically unhelpful in these studies (LR not significant; see [EBM Box 49.1](#)).

Assessment of the patient's traditional risk factors—hypertension, diabetes mellitus, cigarette smoking, family history, or combinations of these—carry much less diagnostic weight than the patient's description of pain. Each of these risk factors—except for cholesterol level higher than 300 mg/dL (LR = 4) and cholesterol level lower than 200 mg/dL (LR = 0.3)—has an LR between the values of 1.2 and 2.3, thus changing probability of disease little if at all.^{79,81,82} Even combinations of three or more risk factors change probability of coronary disease relatively little (LR = 2.2, a value similar to the LR for the earlobe crease).⁷⁹



EBM BOX 49.2
*Myocardial Infarction**

| Finding (Reference) [†] | Sensitivity (%) | Specificity (%) | Likelihood Ratio [‡] if Finding Is | |
|---|--------------------|--------------------|--|--------|
| | | | Present | Absent |
| Patient Interview | | | | |
| Male sex ⁵⁰⁻⁵⁹ | 59-72 | 24-61 | 1.3 | 0.7 |
| Age ^{50,55,56} | | | | |
| <40 years | 4 | 81 | 0.2 | — |
| 40-59 years | 34 | — | NS | — |
| ≥60 years | 47-74 | 54-68 | 1.5 | — |
| Sharp pain ^{55,60,61} | 8-19 | 59-72 | 0.4 | 1.3 |
| Pleuritic pain ^{55,56,60,61} | 3-19 | 69-82 | 0.3 | 1.2 |
| Positional pain ^{55,56,61} | 3-14 | 75-87 | 0.4 | 1.1 |
| Relief of pain with nitroglycerin ⁶²⁻⁶⁵ | 35-92 | 12-59 | NS | NS |
| Physical Examination | | | | |
| Hand gestures ¹⁸ | | | | |
| Levine sign | 7 | 87 | NS | NS |
| Palm sign | 32 | 63 | NS | NS |
| Arm sign | 18 | 83 | NS | NS |
| Pointing sign | 2 | 95 | NS | NS |
| Chest wall tenderness ^{55,56,60,61} | 3-15 | 64-83 | 0.3 | 1.2 |
| Diaphoretic appearance ^{56,59,60} | 28-56 | 71-94 | 2.2 | 0.7 |
| Pallor ⁵⁹ | 70 | 49 | 1.4 | 0.6 |
| Systolic blood pressure <100 mm Hg ⁵² | 6 | 98 | 3.6 | NS |
| Jugular venous distention ⁵¹ | 10 | 96 | 2.4 | NS |
| Pulmonary crackles ^{51,60} | 20-38 | 82-91 | 2.1 | NS |
| Third heart sound ⁶⁰ | 16 | 95 | 3.2 | NS |
| Electrocardiogram | | | | |
| Normal ^{50,52,55,59,66-69} | 1-13 | 48-77 | 0.2 | 1.5 |
| Nonspecific ST changes ^{55,59,68} | 5-8 | 47-78 | 0.2 | 1.4 |
| ST elevation ^{52,59,60,67,68,70,71} | 31-56 | 97-100 | 22.3 | 0.6 |
| ST depression ^{52,59,60,67,68} | 20-62 | 79-96 | 3.9 | 0.8 |
| T wave inversion ^{52,59,60,67} | 9-39 | 84-94 | 2.0 | NS |

*Diagnostic standard: for *myocardial infarction*, development of new electrocardiographic Q waves, elevations of cardiac biomarkers (CK-MB or troponin), or both; except for the studies of nitroglycerin effect, which used a broader definition of "active coronary disease" that combined myocardial infarction, positive stress test, or abnormal coronary arteriogram.⁶²⁻⁶⁴

[†]Definition of findings: for *relief of pain with nitroglycerin*, nitroglycerin provided moderate or complete relief within. All electrocardiographic abnormalities refer to findings that are new or of unknown duration.

[‡]Likelihood ratio (LR) if finding present = positive LR; LR if finding absent = negative LR
NS, Not significant.

[Click here to access calculator](#)




FIG. 49.1 EARLOBE CREASE. The earlobe crease is a diagonal crease extending from the lowest point on the tragus to the outside of the earlobe. See the text.

B. DIAGNOSING MYOCARDIAL INFARCTION

EBM Box 49.2 summarizes the findings in thousands of patients presenting to emergency departments with sustained acute chest pain unrelated to trauma and unexplained by the chest radiograph. The diagnosis of myocardial infarction was confirmed by the development of new Q waves on the electrocardiogram, elevations of cardiac biomarkers (CK-MB or troponin), or both.

According to the LR_s in Table 45.2, the finding *increasing* the probability of myocardial infarction the most is new electrocardiographic ST elevation (LR = 22.3) or ST depression (LR = 3.9). Several additional physical findings have modest value in diagnosing myocardial infarction: systolic blood pressure lower than 100 mm Hg (LR = 3.6), a third heart sound (LR = 3.2), jugular venous distention (LR = 2.4), diaphoretic appearance (LR = 2.2), and pulmonary crackles (LR = 2.1). Radiation of pain to the right arm (LR = 2.7) increases probability of myocardial infarction more than radiation to the left arm (LR = 1.5).^{50,51,60,61,79,83,84} The only findings *decreasing* the probability of myocardial infarction in these studies are pain that is pleuritic (LR = 0.3), positional (LR = 0.4), or sharp (LR = 0.4); a normal electrocardiogram (LR =

0.2); chest wall tenderness (LR = 0.3); and age younger than 40 years (LR = 0.2). In another study of 1635 patients presenting with chest pain to emergency departments, the finding of chest wall tenderness (reproducing the patient's pain) greatly decreased the probability of acute coronary syndrome (i.e., myocardial infarction or unstable angina) during the next 30 days (LR = 0.1).⁸⁵

The response to nitroglycerin fails to discriminate between cardiac and noncardiac causes of chest pain (LR not significant; see [EBM Box 49.2](#)). This may reflect the temporary nature of most chest pain or perhaps the noncardiac effects of nitroglycerin. Nonetheless, even though the response to nitroglycerin lacks diagnostic value in patients with sustained chest pain, it remains a key element in the definition of typical angina. (See the previous discussion in the section on [Description of Chest Pain](#).)

The different hand signs also lack diagnostic value in studies of patients admitted with chest discomfort (see [EBM Box 49.2](#)).

One interesting contrast between the diagnosis of coronary disease (see [EBM Box 49.1](#)) and myocardial infarction (see [EBM Box 49.2](#)) is that chest wall tenderness decreases the probability of myocardial infarction (LR = 0.3; see [EBM Box 49.2](#)) but lacks diagnostic value when considering coronary artery disease (LR = 0.8; see [EBM Box 49.1](#)). This difference may reflect a higher prevalence of chest wall disorders in patients without disease in the acute chest pain studies.

C. RISK FACTORS AND CORONARY DISEASE

In patients with sustained chest pain, the presence or absence of traditional cardiovascular risk factors again carries little or no diagnostic weight (positive LRs = 1.2 to 1.7).⁷⁹ There are two important reasons why risk factors fail to discriminate well in diagnostic studies. First, traditional cardiovascular risk factors are mostly derived from study of middle-aged white residents of Framingham, Massachusetts.⁸⁶ They may thus overestimate the risk in other populations, something that has been demonstrated in British men,⁸⁷ elderly Americans,⁸⁸ and Japanese-American, Native American, and Hispanic populations.⁸⁹ A second reason is the fundamental difference between risk factors and diagnostic signs. Risk factors precede disease, presumably play a role in causing the disease, and become apparent only after study of large groups of *asymptomatic* individuals for long periods of time. In contrast, diagnostic signs first appear after the onset of disease, are *caused by* the disease, and become evident after study of a relatively smaller group of *symptomatic* individuals. For example, it is possible that certain risk factors associated with coronary disease are also associated with noncardiac causes of pain, which would neutralize any diagnostic value (e.g., cigarette smoking may also increase the risk of chest wall pain, making it appear just as often in patients with noncardiac pain as those with cardiac pain. The resulting LR would therefore have a value near 1.).

D. GI COCKTAIL

The existing literature suggests that the GI cocktail has questionable diagnostic value. One problem is that clinicians usually administer the GI cocktail just minutes away from other active medications, such as narcotics, nitroglycerin, antiemetics, histamine blockers, or ketorolac, thus clouding interpretation of the test's results.⁹⁰ Another problem is that the viscous lidocaine is absorbed, and even though most patients have levels below 1 µg/mL (usual therapeutic levels are 2 to 5 µg/mL), instances of toxicity and seizures have occurred.⁹⁰⁻⁹² A final and most troubling problem is the many documented examples of GI cocktail relieving the discomfort of disorders distant from the gastroesophageal mucosa, such as myocardial infarction,^{91,93} hepatitis, pancreatitis, or cholecystitis.⁹⁴

E. PROGNOSIS AND ACUTE CHEST PAIN

In patients with acute chest pain, clinicians are interested in diagnosing more than just myocardial infarction because many acute coronary syndromes *without* infarction are also associated with life-threatening complications that require intensive monitoring and treatment. To identify all patients at risk for such complications, Goldman has developed a rule that assesses the patient's electrocardiogram and the presence or absence of three additional bedside findings: (1) systolic blood pressure of less than 110 mm Hg, (2) crackles heard above the bases bilaterally, and (3) chest pain that is either worse than prior angina, the same as prior myocardial infarction, or occurs in the post-infarction or post-revascularization setting.⁹⁵ According to this rule, patients have a "high risk" of life-threatening complications in the first 24 hours of hospitalization if there is either (1) electrocardiograph ST elevation or Q waves (not known to be old) or (2) electrocardiographic ST depression or T wave inversion (not known to be old) *and* two or more of the three bedside findings. Patients are classified as "very low risk" if their electrocardiogram reveals no ST/T wave changes or Q waves and they lack all three bedside findings.

EBM Box 49.3 indicates that in patients with acute chest pain a "high risk" classification increases the likelihood of life-threatening complications in the subsequent 24 hours (LR = 8.7; see EBM Box 49.3), whereas a "very low risk" classification



EBM BOX 49.3

Predicting Life-Threatening Complications in Patients With Acute Chest Pain*


| Finding (Reference) [†] | Sensitivity (%) | Specificity (%) | Likelihood Ratio [‡] if Finding Is | |
|-------------------------------------|--------------------|--------------------|--|--------|
| | | | Present | Absent |
| Goldman Classification | | | | |
| “High” risk ^{95,96} | 51-88 | 92-93 | 8.7 | — |
| “Very low” risk ⁹⁵⁻⁹⁷ | 7-13 | 42-53 | 0.1 | — |

*Diagnostic standard: for life-threatening complications, any of the following during the first 24 hours of hospitalization: arrhythmias (ventricular fibrillation, cardiac arrest, new complete heart block, insertion of temporary pacemaker, emergency cardioversion), pump failure (cardiogenic shock, use of intra-aortic balloon pump, intubation), or ischemia (recurrent ischemic chest pain requiring bypass surgery or percutaneous intervention).⁹⁵

[†]Definition of findings: for high risk and very low risk, see the text.

[‡]Likelihood ratio (LR) if finding present = positive LR; LR if finding absent = negative LR.

[Click here to access calculator](#)



indicates a favorable prognosis (LR = 0.1; see [EBM Box 49.3](#)). This rule compares favorably with the diagnostic accuracy of elevated troponin T levels, drawn at least 6 hours after the onset of chest pain in patients without ST elevation, in predicting cardiac events in the subsequent 30 days (positive LR = 6.1, negative LR = 0.2).⁹⁸

The references for this chapter can be found on www.expertconsult.com.

REFERENCES

1. Heron M, Hoyert DL, Murphy SL, Xu J, Kochanek KD, Tejada-Vera B. Deaths: final data for 2006. *Natl Vital Stat Rep.* 2006;57(14).
2. Kroenke K, Mangelsdorf AD. Common symptoms in ambulatory care: incidence, evaluation, therapy, and outcome. *Am J Med.* 1989;86:262–266.
3. Qamar A, McPherson C, Babb J, et al. The Goldman algorithm revisited: prospective evaluation of a computer-derived algorithm versus unaided physician judgment in suspected acute myocardial infarction. *Am Heart J.* 1999;138:705–709.
4. Burt CW. Summary statistics for acute cardiac ischemia and chest pain visits to United States EDs, 1995–1996. *Am J Emerg Med.* 1999;17:552–559.
5. Lee TH, Rouan GW, Weisberg MC. Clinical characteristics and natural history of patients with acute myocardial infarction sent home from the emergency room. *Am J Cardiol.* 1987;60:219–224.
6. Goldman L, Weinberg M, Weisberg M, et al. A computer-derived protocol to aid the diagnosis of emergency room patients with acute chest pain. *N Engl J Med.* 1982;307:588–596.
7. Pozen MW, D'Agostino RB, Selker HP, Sytkowski PA, Hood WB. A predictive instrument to improve coronary-care-unit admission practices in acute ischemic heart disease: a prospective multicenter clinical trial. *N Engl J Med.* 1984;310:1273–1278.
8. Pope JH, Aufderheide TP, Ruthazer R, et al. Missed diagnoses of acute cardiac ischemia in the emergency department. *N Engl J Med.* 2000;342:1163–1170.
9. McCarthy BD, Beshansky JR, D'Agostino RB, Selker HP. Missed diagnoses of acute myocardial infarction in the emergency department: results from a multicenter study. *Ann Emerg Med.* 1993;22(3):579–582.
10. Schor S, Behar S, Modan B, Barell V, Drory J, Kariv I. Disposition of presumed coronary patients from an emergency room: a follow-up study. *J Am Med Assoc.* 1976;236:941–943.
11. Schull MJ, Vermeulen MJ, Stukel TA. The risk of missed diagnosis of acute myocardial infarction associated with emergency department volume. *Ann Emerg Med.* 2006;48:647–655.
12. Collinson PO, Premachandram S, Hashemi K. Prospective audit of incidence of prognostically important myocardial damage in patients discharged from emergency department. *Br Med J.* 2000;320:1702–1705.
13. Osler W. *Lectures on Angina Pectoris and Allied States.* 1st ed. New York, NY: Appleton & Co.; 1897.
14. Major R. *Classic Descriptions of Disease: with Biographical Sketches of the Authors.* Springfield, IL: Charles C Thomas; 1932.
15. Burch GE, DePasquale NP. *A History of Electrocardiography.* Chicago, IL: Year Book Medical Publishers, Inc; 1964.
16. Flint A. *A Practical Treatise on the Diagnosis, Pathology, and Treatment of Diseases of the Heart.* Philadelphia, PA: Blanchard and Lea; 1859:260.
17. Eslick GD. Chest pain: a historical perspective. *Int J Cardiol.* 2001;77:5–11.
18. Marcus GM, Cohen J, Varosy PD, et al. The utility of gestures in patients with chest discomfort. *Am J Med.* 2007;120:83–89.
19. Weiner DA, Ryan TJ, McCabe CH, et al. Exercise stress testing: correlations among history of angina, ST-segment response and prevalence of coronary-artery disease in the coronary artery surgery study (CASS). *N Engl J Med.* 1979;301:230–235.
20. Proudfit WL, Shirey EK. Selective cine coronary arteriography: correlation with clinical findings in 1000 patients. *Circulation.* 1966;33:901–910.
21. Peissens J, van Mieghem W, Kesteloot H, de Geest H. Diagnostic value of clinical history, exercise testing and atrial pacing in patients with chest pain. *Am J Cardiol.* 1974;33:351–356.
22. McConahay DR, McCallister BD, Smith RE. Postexercise electrocardiography: correlations with coronary arteriography and left ventricular hemodynamics. *Am J Cardiol.* 1971;28:1–9.
23. Mason RE, Likar I, Biern RO, Ross RS. Multiple-lead exercise electrocardiography: experience in 107 normal subjects and 67 patients with angina pectoris, and comparison with coronary cinearteriography in 84 patients. *Circulation.* 1967;34:517–525.
24. Detry JMR, Kapita BM, Cosyns J, Sottiaux B, Brasseur LA, Rousseau MF. Diagnostic value of history and maximal exercise electrocardiography in men and women suspected of coronary heart disease. *Circulation.* 1977;56:756–769.

25. Chaitman BR, Bourassa MG, Davis K, et al. Angiographic prevalence of high-risk coronary artery disease in patient subsets (CASS). *Circulation*. 1981;64:360–367.
26. Campeau L, Bourassa MG, Bois MA, et al. Clinical significance of selective coronary cinearteriography. *Can Med Assoc J*. 1968;99:1063–1068.
27. Cooke RA, Smeeton N, Chambers JB. Comparative study of chest pain characteristics in patients with normal and abnormal coronary angiograms. *Heart*. 1997;78:142–146.
28. Pearson TA, Bulkley BH, Achuff SC, Kwiterovich PO, Gordis L. The association of low levels of HDL cholesterol and arteriographically defined coronary artery disease. *Am J Epidemiol*. 1979;109(3):285–295.
29. Toyosaki N, Tsuchiya M, Hashimoto T, et al. Earlobe crease and coronary heart disease in Japanese. *Heart Vessels*. 1986;2:161–165.
30. Holmes DR, Elveback LR, Frye RL, Kottke BA, Ellefson RD. Association of risk factor variables and coronary artery disease documented with angiography. *Circulation*. 1981;63(2):293–299.
31. Hartz A, Grubb B, Wild R, et al. The association of waist hip ratio and angiographically determined coronary artery disease. *Int J Obes*. 1990;14:657–665.
32. Haft JI, Gonnella GR, Kirtane JS, Anastasiades A. Correlation of ear crease sign with coronary arteriographic findings. *Cardiovasc Med*. 1979;4:861–867.
33. Pryor DB, Shaw L, Harrell FE, et al. Estimating the likelihood of severe coronary artery disease. *Am J Med*. 1991;80:553–562.
34. Elder AT, Shaw TRD, Turnbull CM, Starkey IR. Elderly and younger patients selected to undergo coronary angiography. *Br Med J*. 1991;303:950–953.
35. Gurevitz O, Janos M, Boyko V, Rabinowitz B, Reicher-Reiss H. Clinical profile and long-term prognosis of women <50 years of age referred for coronary angiography for evaluation of chest pain. *Am J Cardiol*. 2000;85:806–809.
36. Miranda CP, Lehmann KG, Froelicher VF. Correlation between resting ST segment depression, exercise testing, coronary angiography, and long-term prognosis. *Am Heart J*. 1991;122:1617–1628.
37. Elliott WJ, Powell LH. Diagonal earlobe creases and prognosis in patients with suspected coronary artery disease. *Am J Med*. 1996;100:205–211.
38. Brady PM, Zive MA, Goldberg RJ, Gore JM, Dalen JE. A new wrinkle to the earlobe crease. *Arch Intern Med*. 1987;147:65–66.
39. Kaukola S, Manninen V, Valle M, Halonen PI. Ear-lobe crease and coronary atherosclerosis. *Lancet*. 1979;2:1377.
40. Wu XL, Yag D, Zhao Y, Jin ML. Diagonal earlobe crease and coronary artery disease in a Chinese population. *BMC Cardiovasc Disord*. 2014;14:43.
41. Evrengül H, Dursunoglu D, Kaftan A, et al. Bilateral diagonal earlobe crease and coronary artery disease: a significant association. *Dermatology*. 2004;209:271–275.
42. Hoogerbrugge N, Happee C, va Domburg R, Poldermans D, van der Brand MJB. Corneal arcus: indicator for severity of coronary atherosclerosis? *Neth J Med*. 1999;55:184–187.
43. Levine PR, Mascette AM. Musculoskeletal chest pain in patients with “angina”: a prospective study. *South Med J*. 1989;82(5):580–585.
44. Wu EB, Smeeton N, Chambers JB. A chest pain score for stratifying the risk of coronary artery disease in patients having day case coronary angiography. *Int J Cardiol*. 2001;78:257–264.
45. Kumarathurai P, Farooq MK, Christensen HW, Vach W, Hoiland-Carlson PF. Muscular tenderness in the anterior chest wall in patients with stable angina pectoris is associated with normal myocardial perfusion. *J Manipulative Physiol Ther*. 2008;31:344–347.
46. Papamichael CM, Lekakis JP, Stamatelopoulos KS, et al. Ankle-brachial index as a predictor of the extent of coronary atherosclerosis and cardiovascular events in patients with coronary artery disease. *Am J Cardiol*. 2000;86:615–618.
47. Chang ST, Chen CL, Chu CM, et al. Ankle-arm index is a useful test for clinical practice in outpatients with suspected coronary artery disease. *Circ J*. 2006;70:686–690.
48. Mukerji V, Alpert MA, Hewett JE, Parker BM. Can patients with chest pain and normal coronary arteries be discriminated from those with coronary artery disease prior to coronary angiography? *Angiology*. 1989;40:276–282.
49. Linhart JW, Turnoff HB. Maximum treadmill exercise test in patients with abnormal control electrocardiograms. *Circulation*. 1974;49:667–672.

50. Berger JP, Buclin T, Haller E, Van Melle G, Yersin B. Right arm involvement and pain extension can help to differentiate coronary diseases from chest pain of other origin: a prospective emergency ward study of 278 consecutive patients admitted for chest pain. *J Intern Med.* 1990;227:165–172.
51. Baxt WG. Use of an artificial neural network for the diagnosis of myocardial infarction. *Ann Intern Med.* 1991;115:843–848.
52. Herlitz J, Karlson BW, Karlsson T, Svensson L, Kalin EZB. A description of the characteristics and outcome of patients hospitalized for acute chest pain in relation to whether they were admitted to the coronary care unit or not in the thrombolytic era. *Int J Cardiol.* 2002;82:279–287.
53. Herlitz J, Starke M, Hansson E, Ringvall E, Karlson BW, Waagstein L. Early identification of patients with an acute coronary syndrome as assessed by dispatchers and the ambulance crew. *Am J Emerg Med.* 2002;20(3):196–201.
54. Herlitz J, Bang A, Isaksson L, Karlsson T. Ambulance dispatchers' estimation of intensity of pain and presence of associated symptoms in relation to outcome in patients who call for an ambulance because of acute chest pain. *Eur Heart J.* 1995;16:1789–1794.
55. Lee TH, Cook EF, Weisberg M, Sargent RK, Wilson C, Goldman L. Acute chest pain in the emergency room: identification and examination of low-risk patients. *Arch Intern Med.* 1985;145:65–69.
56. Solomon CG, Lee TH, Cook EF, et al. Comparison of clinical presentation of acute myocardial infarction in patients older than 65 years of age to younger patients: the multicenter chest pain study experience. *Am J Cardiol.* 1989;63:772–776.
57. Bansal D, Eigenbrodt M, Gupta E, Mehta JL. Traditional risk factors and acute myocardial infarction in patients hospitalized with cocaine-associated chest pain. *Clin Cardiol.* 2007;30:290–294.
58. Body R, McDowell G, Carley S, Mackway-Jones K. Do risk factors for chronic coronary heart disease help diagnose acute myocardial infarction in the emergency department? *Resuscitation.* 2008;79:41–45.
59. Svensson L, Isaksson L, Axelsson C, Nordlander R, Herlitz J. Predictors of myocardial damage prior to hospital admission among patients with acute chest pain or other symptoms raising a suspicion of acute coronary syndrome. *Coron Artery Dis.* 2003;14:225–231.
60. Tierney WM, Fitzgerald J, McHenry R, et al. Physicians' estimates of the probability of myocardial infarction in emergency room patients with chest pain. *Med Decis Making.* 1986;6:12–17.
61. Rubini Gimenez M, Reiter M, Twerenbold R, et al. Sex-specific chest pain characteristics in the early diagnosis of acute myocardial infarction. *JAMA Intern Med.* 2013;174:241–249.
62. Shry EA, Dacus J, Van de Graaff E, Hjelkrem M, Stajduhar KC, Steinhilber SR. Usefulness of the response to sublingual nitroglycerin as a predictor of ischemic chest pain in the emergency department. *Am J Cardiol.* 2002;90:1264–1266.
63. Henrikson CA, Howell EE, Bush DE, et al. Chest pain relief by nitroglycerin does not predict active coronary artery disease. *Ann Intern Med.* 2003;139:979–986.
64. Diercks DB, Boghes E, Guzman H, Amsterdam EA, Kirk JD. Changes in the numeric descriptive scale for pain after sublingual nitroglycerin do not predict cardiac etiology of chest pain. *Ann Emerg Med.* 2005;45(6):581–585.
65. Steele R, McNaughton T, McConahy M, Lam J. Chest pain in emergency department patients: if the pain is relieved by nitroglycerin, is it more likely to be cardiac chest pain? *Can Assoc Emerg Phys.* 2006;8(3):164–169.
66. Brush JE, Brand DA, Acampora D, Chalmer B, Wackers FJ. Use of the initial electrocardiogram to predict in-hospital complications of acute myocardial infarction. *N Engl J Med.* 1985;312:1137–1141.
67. Karlson BW, Herlitz J, Wiklund O, Richter A, Hjalmarson A. Early prediction of acute myocardial infarction from clinical history, examination and electrocardiogram in the emergency room. *Am J Cardiol.* 1991;68:171–175.
68. Rouan GW, Lee TH, Cook EF, Brand DA, Weisberg MC, Goldman L. Clinical characteristics and outcome of acute myocardial infarction in patients with initially normal or nonspecific electrocardiograms (A report from the Multicenter Chest Pain Study). *Am J Cardiol.* 1989;64:1087–1092.

69. Challa PK, Smith KM, Conti CR. Initial presenting electrocardiogram as determinant for hospital admission in patients presenting to the emergency department with chest pain: a pilot investigation. *Clin Cardiol.* 2007;30:558–561.
70. Aufderheide TP, Hendley GE, Woo J, Lawrence S, Valley V, Teichman SL. A prospective evaluation of prehospital 12-lead ECG application in chest pain patients. *J Electrocardiol.* 1992;24(suppl):8–13.
71. Fesmire FM, Percy RF, Bardoner JB, Wharton DR, Calhoun FB. Usefulness of automated serial 12-lead ECG monitoring during the initial emergency department evaluation of patients with chest pain. *Ann Emerg Med.* 1998;31:3–11.
72. Frank ST. Aural sign of coronary-artery disease. *N Engl J Med.* 1973;289:327–328.
73. Elliott WJ, Karrison T. Increased all-cause and cardiac morbidity and mortality associated with the diagonal earlobe crease: a prospective cohort study. *Am J Med.* 1991;91:247–254.
74. Ishii T, Asuwa N, Masuda S, Ishikawa Y, Shimada K, Takemoto S. Earlobe crease and atherosclerosis: an autopsy study. *J Am Geriatr Soc.* 1990;38:871–876.
75. McAndrew GM, Ogston D. Arcus senilis and coronary artery disease. *Am Heart J.* 1965;70(6):838–840.
76. Chambless LE, Fuchs FD, Linn S, et al. The association of corneal arcus with coronary heart disease and cardiovascular disease mortality in the Lipid Research Clinics mortality follow-up study. *Am J Public Health.* 1990;80:1200–1204.
77. Rosenman RH, Brand RJ, Sholtz RI, Jenkins CD. Relation of corneal arcus to cardiovascular risk factors and the incidence of coronary disease. *N Engl J Med.* 1974;291:1322–1324.
78. Berman D, Graber M, Clark P, et al. The GI cocktail in the treatment of dyspepsia: a randomized, double-blind clinical trial. *Acad Emerg Med.* 2001;8:446.
79. Chun AA, McGee SR. Bedside diagnosis of coronary artery disease: a systematic review. *Am J Med.* 2004;117:334–343.
80. Edmondstone WM. Cardiac chest pain: does body language help the diagnosis? *Br Med J.* 1995;311:1660–1661.
81. Welch CC, Proudfit WL, Sheldon WC. Coronary arteriographic findings in 1000 women under age 50. *Am J Cardiol.* 1975;35:211–215.
82. Welch CC, Proudfit WL, Sones FM, Shirey EK, Sheldon WC, Razavi M. Cinecoronary arteriography in young men. *Circulation.* 1970;42:647–652.
83. Albarran J, Durham B, Gowers J, Dwight J, Chappell G. Is the radiation of chest pain a useful indicator of myocardial infarction? A prospective study of 541 patients. *Accident Emerg Nurs.* 2002;10:2–9.
84. Säwe U. Pain in myocardial infarction. *Acta Med Scand.* 1971;190:79–81.
85. Cullen L, Greenslade JH, Than M, et al. The new Vancouver chest pain rule using troponin as the only biomarker: an external validation study. *Am J Emerg Med.* 2013;32:129–134.
86. Truett J, Cornfield J, Kannel W. A multivariate analysis of the risk of coronary heart disease in Framingham. *J Chron Dis.* 1967;20:511–524.
87. Prindle P, Emberson J, Lampe F, et al. Predictive accuracy of the Framingham coronary risk score in British men: prospective cohort study. *Br Med J.* 2003;327:1267–1270.
88. Psaty BM, Anderson M, Kronmal RA, et al. The association between lipid levels and the risks of incident myocardial infarction, stroke, and total mortality: the cardiovascular health study. *J Am Geriatr Soc.* 2004;52:1639–1647.
89. D'Agostino RB, Grundy S, Sullivan LM, Wilson P. Validation of the Framingham coronary heart disease prediction scores: results of a multiple ethnic groups investigation. *JAMA.* 2001;286(2):180–187.
90. Wrenn K, Slovis CM, Gongaware J. Using the “GI cocktail”: a descriptive study. *Ann Emerg Med.* 1995;26:687–690.
91. Schwartz GR. Xylocaine viscous as an aid in the differential diagnosis of chest pain. *JACEP.* 1976;5:981–983.
92. Hess GP, Walson PD. Seizures secondary to oral viscous lidocaine. *Ann Emerg Med.* 1988;17:725–727.
93. Dickinson MW. The GI cocktail in the evaluation of chest pain in the emergency department. *J Emerg Med.* 1996;14:245–246.
94. Welling LR, Watson WA. The emergency department treatment of dyspepsia with antacids and oral lidocaine. *Ann Emerg Med.* 1990;19:785–788.

95. Goldman L, Cook EF, Johnson PA, Brand DA, Rouan GW, Lee TH. Prediction of the need for intensive care in patients who come to emergency departments with acute chest pain. *N Engl J Med*. 1996;334:1498–1504.
96. Reilly B, Durairaj L, Husain S, et al. Performance and potential impact of a chest pain prediction rule in a large public hospital. *Am J Med*. 1999;106:285–291.
97. Durairaj L, Reilly B, Das K, et al. Emergency department admissions to inpatient cardiac telemetry beds: a prospective cohort study of risk stratification and outcomes. *Am J Med*. 2001;110:7–11.
98. Hamm CW, Goldmann BU, Heeschen C, Kreyman G, Berger J, Meinertz T. Emergency room triage of patients with acute chest pain by means of rapid testing for cardiac troponin T or troponin I. *N Engl J Med*. 1997;337:1648–1653.